



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Direct manipulation of T lymphocytes by proteins of gastrointestinal bacterial pathogens

Citation for published version:

Cassady-Cain, RL, Hope, JC & Stevens, MP 2018, 'Direct manipulation of T lymphocytes by proteins of gastrointestinal bacterial pathogens', *Infection and Immunity*, vol. 86, no. 5, 00683-17.
<https://doi.org/10.1128/IAI.00683-17>

Digital Object Identifier (DOI):

[10.1128/IAI.00683-17](https://doi.org/10.1128/IAI.00683-17)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Infection and Immunity

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



1 **Direct manipulation of T lymphocytes by proteins of gastrointestinal bacterial pathogens**

2

3 Robin L. Cassady-Cain*, Jayne C. Hope and Mark P. Stevens

4

5 The Roslin Institute and Royal (Dick) School of Veterinary Studies, University of Edinburgh,

6 Midlothian, EH25 9RG, United Kingdom

7 *Corresponding author

8 Email: robin.cassady-cain@roslin.ed.ac.uk

9 Tel: +44 (0)131 6519238

10

11 **Abstract**

12 Gastrointestinal bacterial infection represents a significant threat to human health, as well
13 as a burden on food animal production and welfare. Although there is advanced knowledge
14 about the molecular mechanisms underlying pathogenesis, including the development of
15 immune responses to these pathogens, gaps in knowledge persist. It is well established that
16 gastrointestinal bacterial pathogens produce a myriad of proteins that affect the
17 development and effectiveness of innate immune responses. However, relatively few
18 proteins that directly affect lymphocytes responsible for humoral or cell-mediated immunity
19 and memory have been identified. Here, we review factors produced by gastrointestinal
20 bacterial pathogens that have direct T cell interactions and what is known about their
21 functions and mechanisms of action. T cell interacting bacterial proteins that have been
22 identified to date mainly target three major T cell responses: activation and expansion,
23 chemotaxis or apoptosis. Further, the requirement for more focused studies to identify and
24 understand additional mechanisms used by bacteria to directly affect the T cell immune
25 response and how these may contribute to pathogenesis is highlighted. Increased
26 knowledge in this area will help to drive development of better interventions in prevention
27 and treatment of gastrointestinal bacterial infection.

28

29 Introduction

30 Gastrointestinal bacterial infection represents a significant threat to human health
31 and welfare, with an estimated 900 million illnesses resulting in over 500 000 deaths in a
32 single year, according to the World Health Organisation (1). Although much is known about
33 the molecular mechanisms underlying persistence, pathogenesis and protection, significant
34 effort is still required to devise effective intervention strategies. Bacterial immune evasion
35 methods include expression of surface polysaccharides to resist complement-mediated
36 killing and opsonisation, enzymes to detoxify reactive oxygen species in phagosomes,
37 escape from phagosomes and in the case of intracellular bacterial pathogens, interference
38 in cellular antigen-presentation and innate immune responses by proteins secreted by Type
39 III or Type IV secretion systems (2). The role of Type III secretion in gastrointestinal bacterial
40 pathogens is covered by many high quality reviews (eg. 3), and therefore is not the focus of
41 this review. Relatively fewer mechanisms have been identified whereby bacteria are able to
42 directly affect lymphocytes during infection. Further, understanding the full role of these
43 bacterial proteins and their T cell interactions during infections, any specificity for T cells
44 subsets, and proof that they are able to directly meet T cells in the body is crucial to
45 establish biological importance. However, such evidence is often lacking, and a number of
46 barriers to understanding these aspects exist.

47 Here, we review the known molecules and strategies that contribute to the direct
48 subversion or dampening of the adaptive T cell response in gastrointestinal bacterial
49 infection. In addition, we discuss the challenges and aspirations of identifying these
50 mechanisms.

51

52 T cell distribution in the intestine

53 An understanding of the importance of T-cell targeting strategies by bacteria that
54 infect the intestine requires an understanding of the lymphoid architecture, distribution of
55 gut resident T cells, the ability of T cells to be recruited to this tissue and the nature of the
56 downstream immune response that is triggered upon infection. T cell responses in the
57 intestine are mainly governed by the gut-associated lymphoid tissue (GALT), which is similar
58 to other secondary lymphoid tissues in the body, and in mammals, consists of the
59 mesenteric lymph nodes (mLNs) and Peyer's patches (PP), the appendix and multiple
60 smaller isolated lymphoid follicles studding the intestinal wall. In addition, there are isolated
61 immune cells scattered in the lamina propria (LP) and throughout the epithelium of the
62 intestine (reviewed in 4) that contribute to intestinal immune responses (summarized in
63 Figure 1). T cell responses in the gut can be initiated from several sites. Generally in an
64 intestinal immune response, intravascular naïve T cells home to the GALT (specifically to the
65 PP and mLN), where they can meet their cognate ligand in the context of the Major
66 Histocompatibility Complex (MHC) and become activated. These T cells are then able to exit
67 the lymphoid tissue via the lymphatic vessels and enter the circulation to home back to the
68 intestinal LP (reviewed in 5) where they are able to carry out their functions. Within the
69 mucosa, dendritic cells in the LP sample antigens and migrate to the PP and mLN where they
70 are able to prime and present antigen to naïve CD4⁺ and CD8⁺ T cells, which clonally
71 expand. These T cells may become memory cells, which accumulate over time in the LP. The
72 LP is mainly enriched for the CD4⁺ T_{reg} and T_h17 cells. In contrast, the intraepithelial
73 lymphocyte (IEL) resident T cell population is mainly composed of both T cell receptor (TCR)
74 αβ and TCR γδ (TCR αβ are generally considered "conventional T cells", TCR γδ cells are
75 often considered non-conventional atypical T cells), both mainly CD8ββ isoform. However,
76 IELs appear to lack some typical T cell surface molecules such as CD2 (adhesion molecule),

77 CD28 (activating co-receptor) and Thy-1 (pan T cell marker of human and mouse cells).
78 These cells are considered “activated, yet resting” and are different compared to peripheral
79 T cells (which express CD4 or CD8 $\alpha\beta$ isoform; reviewed in (4)).

80 Although there is not a large literature describing direct effects of gastrointestinal
81 bacterial pathogens on T cells, a picture of the general strategies used to alter T lymphocyte
82 function is emerging (summarised in Figure 2). The effects identified can be broadly
83 assigned to three groups: those that affect T cell activation and proliferation, those that
84 affect chemotaxis and those that cause elimination of T cells, with most of the strategies
85 identified thus far falling into the first category. Little duplication of specific strategies
86 across species of intestinal bacteria has been identified so far, and the mechanisms and
87 observations described are derived from a relatively small number of bacterial species.

88 **Proteins that affect T cell activation and proliferation**

89 A number of bacterial proteins have been identified that act to interfere with signalling
90 cascades in T cell activation and expansion, which are outlined below. These proteins are
91 often soluble, diffusible factors, and can act externally to the T cell, as well as intracellularly.

92 *Superantigens*

93 Perhaps the earliest, well known and characterised bacterial factor with T cell-
94 affecting activity is superantigen. Barber observed superantigen activity in *Staphylococcus*
95 in 1914, and identified the cause to be a microorganism-derived toxin (6). Since then, there
96 have been numerous studies characterising the activity of superantigens. One of the
97 hallmarks of superantigens is their ability to activate a large population of T cells at very low
98 concentrations (7, 8). At a basic level, superantigens are able to cross-link a relatively large
99 number of T cells to antigen-presenting cells (APC) compared to normal antigen-driven
100 activation, inducing wide spread non-antigen-specific activation of T cells, ultimately leading

101 to clonal deletion and anergy, thus suppressing a productive T cell response (9).

102 Superantigens are effective because they bind outside the peptide-binding groove of MHC,
103 they are not MHC restricted and activation does not rely on antigen internalisation and
104 processing. In addition, they specifically require the TCR β chain, not the $V\alpha$ - $V\beta$ chain
105 pairing required in conventional antigen recognition by the TCR (reviewed in 8).

106 Although most well described in *Staphylococcus*, superantigens have been identified
107 in other bacteria, including the pathogens *Yersinia enterocolitica* (10), and *Yersinia*
108 *pseudotuberculosis* (11), the latter of which most often causes a self-limiting gastrointestinal
109 infection. However, strains of *Y. pseudotuberculosis* have also been reported to infect the
110 gut and cause Far East Scarlet-like Fever (reviewed in 16), and many strains associated with
111 this pathogenic infection express the superantigen *Y. pseudotuberculosis*-derived mitogen A
112 (YPMa). Strains deficient in YPMa have been demonstrated to have decreased
113 pathogenicity, however, growth of the bacteria was unaffected in the major immune organs
114 after oral infection, so YPMa may have more pronounced effects in systemic infection (13),
115 and a more recent study has linked the toxic activity of YPMa to activation of a hepatotoxic
116 $CD4^+$ T cell subset (14).

117 *Lymphostatin*

118 Lymphostatin (LifA, Efa-1) is one of the largest known bacterial proteins at 365 kDa,
119 and is a putative glycosyltransferase, expressed by enteropathogenic *Escherichia coli* (EPEC)
120 and non-O157 enterohaemorrhagic *E. coli* (EHEC) (15). It has homology to the large
121 clostridial toxins A and B (TcdA/B) at the N terminal portion of the protein, where the
122 catalytic glycosyltransferase domain of TcdA/B resides (15, 16). The existence of a soluble
123 factor capable of inhibiting mitogen-activated lymphocyte proliferation and pro-
124 inflammatory cytokine expression was first described using crude bacterial lysates of the

125 prototype EPEC strain E3248/69 (17). Lymphostatin was then subsequently identified using
126 a cosmid library screen to identify the gene responsible for this activity, which was
127 confirmed by mutation of the *lifA* gene in EPEC (15). Recently, it was shown that
128 lymphostatin, like its clostridial homologues, is able to bind sugar moieties, in this case UDP-
129 N-acetyl glucosamine (UDP-GlcNAc), and has significant predicted structural homology
130 around the putative glycosyltransferase domain. A DXD motif within this domain is
131 necessary for UDP-GlcNAc binding and lymphostatin activity, however, formal evidence of
132 sugar transfer and the identity of the cellular target remain elusive (16). It has been
133 demonstrated that lymphostatin is capable of inhibiting all major T cell subsets. In addition,
134 lymphostatin has some activity against B cells, but not natural killer cells (18). Further, the
135 effects of lymphostatin on T cells appear to be long-lived, even in the absence of continued
136 incubation with the protein, preventing mitogenic activation for more than 18 hours after
137 transient exposure and withdrawal of the protein. Lymphostatin was also able to inhibit
138 antigen-specific proliferation of bovine T cells using *Theileria parva* antigens presented on
139 infected irradiated APCs to *T. parva* specific T cells as a model antigen system (18). These
140 findings suggest that lymphostatin might act to permanently de-sensitise T cells to stimulus,
141 possibly suppressing T cell responses and preventing or dampening a productive immune
142 response and delaying clearance of infection (18). It would appear that the effects of
143 lymphostatin interfere with signalling in a membrane proximal way, as inhibition was not
144 achieved in T cells stimulated with Phorbol 12-myristate 13-acetate (PMA)/ionomycin,
145 which bypass membrane signalling.

146 Lymphostatin is known to play an important role in intestinal colonization of calves
147 by non-O157 EHEC strains of multiple serogroups (19, 20) and of mice by *Citrobacter*
148 *rodentium* (21). However, attenuation is evident early after infection, before adaptive

149 responses may be expected to have developed. Alongside the ability to suppress T cell
150 activation, lymphostatin also appears to be associated with adhesion (22), possibly as a
151 consequence of effects on Type III secretion in some strains (19, 20). These results indicate
152 that lymphostatin may have additional roles in infection. There are a number of unresolved
153 questions regarding the activity of lymphostatin, including its cellular target of glycosylation.
154 Further, in O157:H7 strains of EHEC, where full-length lymphostatin is not expressed, there
155 is a putative homologue, ToxB, that also shows T cell inhibitory activity, as well as homology
156 at the N-terminal end of the molecule to TcdA/B (18). This suggests that lymphostatin and
157 lymphostatin-like molecules may be a family of proteins expressed by *E. coli* to control T cell
158 responses to infection.

159

160 *VacA*

161 *Helicobacter pylori* expresses the *VacA* vacuolating cytotoxin, which has direct
162 activity against T cells, specifically inhibiting T cell proliferation (23–25) as well as effects on
163 other cells, including phagocytes and epithelial cells (likely by a different mechanism;
164 reviewed in 23). *VacA* is a two domain protein, processed from a protoxin form, after
165 secretion via a Type Va system from the bacteria (reviewed in 24). Variation in the *VacA*
166 gene amongst different strains of *H. pylori* results in varying levels of toxicity among the
167 different variants (28). Like other toxins, *VacA* must be taken up by the cell in order to exert
168 its activity, and it has been shown that both domains are needed for proper uptake and
169 function of the toxin (29). The integrin CD18, expressed on the cell surface, has been
170 identified as being important for uptake of *VacA* in human T cells (30), mediated by Protein
171 Kinase C (PKC), and activation of the T cell is required to see the active endocytosis of *VacA*
172 in T cells (31). In addition, *VacA* is able to block calcium flux in the Jurkat T cell line (32), and

173 prevent IL-2 expression by blocking translocation of the transcription factor NF-AT (24, 25).

174 Overall, the data suggest that VacA targets previously activated T cells. Using *in vivo* studies
175 in mice, a null mutation of *vacA* was reported to impair initial colonization of mice by *H.*
176 *pylori*, however, once infection by the *vacA* mutant becomes established, the bacterial load
177 and extent of intestinal inflammation were similar to the parent strain (33). This effect is
178 independent of an effect on T cells, as mice T cells do not express a compatible receptor
179 that allows uptake of VacA (30).

180

181 *YopH*

182 Another example of inhibition of T cell activation by interference in T cell signalling is
183 the YopH protein from *Yersinia*. YopH is expressed by *Yersinia spp.* that infect the gut (34),
184 including *Y. enterocolitica* and *Y. pseudotuberculosis*, and has been characterised as a protein
185 tyrosine phosphatase (35). In *in vitro* studies, using T cell-like cell lines, YopH was able to
186 inhibit IL-2 production induced by antigen stimulation, the effects of which were upstream
187 of PMA/Ionomycin (36). It was apparent by Western Blotting that general tyrosine
188 phosphorylation of signalling molecules was inhibited. YopH has also been shown to exhibit
189 activity against B cell activation via the B cell receptor, with similar characteristics (36).
190 These effects were independently confirmed in primary human T cells (37). In T cells, YopH
191 is able to dephosphorylate the early signalling molecule Lck (38). Further, it has been shown
192 that YopH interacts with a number of adaptor molecules involved in early T cell receptor
193 signalling. Using a trapping mutant, YopH was shown to directly dephosphorylate
194 recombinant phosphorylated Lck in an *in vitro* activity assay, while not
195 dephosphorylating other associated adaptor molecules, indicating some specificity of
196 activity (39). This is an elegant mechanism, as an effect on relatively few molecules of Lck

197 would have a large impact on downstream signalling due to amplification through the
198 signalling cascade. These studies remain quite far removed from the complex *in vivo*
199 infection, so the implications of these activities are not entirely known. However, it has
200 been demonstrated *in vivo* that *yopH* deficient *Y. enterocolita* are drastically attenuated in
201 oral infection of C57BL/6 mice, although colonization of the small intestine persists until at
202 least 21 days post-infection (40). Colonization by the YopH mutant declined quickly after
203 infection (40). Further, in an intranasal infection model, a *yopH* deficient strain was less
204 effective at lung colonization (41). In both cases, reduced colonization was seen early in
205 infection before one might expect an adaptive response to have properly formed, and so it
206 remains unclear what role YopH/T cell interactions play in virulence. In addition to YopH,
207 *Yersinia* expresses an additional protein, invasin, which may allow *Yersinia* to subvert
208 lymphocytes, particularly T cells, to influence their motility and facilitate dissemination of
209 *Yersinia* to distal sites (42). These two proteins may function to simultaneously neutralize T
210 cell activation while keeping the cells intact to allow *Yersinia* infection, and redirection to
211 other sites within the body. An invasin homologue in EPEC and *Citrobacter rodentium*,
212 intimin (reviewed in 41), has been shown to interact with T cells, however, it is difficult to
213 separate its direct effects on lymphocytes from the vital role it plays in gut colonization
214 when interpreting its role *in vivo* (44, 45).

215

216 *Interference with metabolic activity*

217 T cells undergo rapid metabolic reprogramming on activation, one of the
218 requirements of which is a source extracellular amino acids (46). Import of amino acids such
219 as asparagine and glutamine is required to accommodate the increased metabolic load
220 induced by aerobic glycolysis during activation and proliferation of T cells (47). There are at

221 least two examples of proteins from gastrointestinal bacterial pathogens that appear to
222 inhibit T cell activation via limiting availability of extracellular amino acids.

223 In a recent publication, Floch and colleagues (48) reported that the *Campylobacter*
224 *jejuni* protein, gamma-glutamyl transpeptidase (GGT) was capable of inhibiting mitogenic
225 proliferation of T cells *in vitro*. Although GGT is known to be important in intestinal
226 colonization by *C. jejuni* in the chicken (49), little is known about its activity on T cells.
227 However, it is tempting to extrapolate from what is known about a similar GGT that is
228 expressed by *H. pylori*. The GGT of *H. pylori* plays an essential role in colonization of the
229 gastric mucosa in mice (50). GGTs are N-terminal nucleophile hydrolases that play a role in
230 the degradation of glutathione, and GGTs across mammal and bacterial species often
231 exhibit a high protein sequence identity, with the GGT of *C. jejuni* clustering with
232 *Helicobacter spp.* (51). Treatment of mouse T cells with recombinant GGT from *H. suis*
233 inhibits CD3/CD28-stimulated proliferation in a concentration-dependent manner (52). In
234 human peripheral blood mononuclear cells, GGT also inhibits PMA/Ionomycin stimulated
235 proliferation, causing cell cycle arrest, inhibiting c-Myc and c-Raf (53). GGT more specifically
236 causes glutamine deprivation in the extracellular space of T cells, downregulating both c-
237 Myc and IRF4 which are sensitive to glutamine, and required for metabolic adaptation (54).
238 Overall, the data suggest that GGT is able to modulate the response of T cells in infection,
239 likely through control of the extracellular availability of glutamine, which is required during
240 activation.

241 A second example of a gastrointestinal bacterial protein that interferes with T cell
242 metabolism comes from *Salmonella enterica* serovar Typhimurium, which has been
243 reported to directly inhibit primary mouse T cells (55), and is thought to limit availability of
244 asparagine to T cells (56). When assessing a number of cell surface expressed molecules, no

245 difference was noted in levels of CD69, CD25 α , CD44, and CD62L in cells infected with *S.*
246 Typhimurium compared to uninfected controls. However, in the presence of *S.*
247 Typhimurium, neither IL-2 nor IFN- γ were produced with CD3 cross-linking (both these
248 cytokines are up-regulated during T cell activation). Cytokine production was restored if the
249 cells were separated from the bacteria in a transwell arrangement, indicating that direct
250 contact with the bacteria was required for this effect. From here, the authors extended
251 their observations, again using *in vitro* methods with primary mouse cells, showing that *S.*
252 Typhimurium was able to down regulate surface expression the TCR β chain, resulting in
253 decreased gene expression, intracellular and surface protein, at least partially explaining the
254 mechanism targeted to inhibit T cell activation (57). Not only that, but this effect was only
255 observed in the presence of live bacteria, as treatment with heat-inactivated bacteria
256 abrogated this effect. The effects were shown to be unrelated to Type III secretion or the
257 bacterial virulence plasmid. It was later shown that the protein responsible for this was L-
258 asparaginase II (STM3106; *asnB*) (58). In a mouse model of bacterial persistence, the burden
259 of bacteria was lower in mutants lacking L-asparaginase II, suggesting that this molecule
260 may enable bacterial persistence by dampening the T cell-mediated immune response (58).
261 In contrast, in a screen of *S. Typhimurium* mutants in pigs, calves and chickens, a transposon
262 insertion was not attenuating in the gut, albeit within 3-4 days after oral infection (59).
263 Nonetheless, the characterization of the activity of the L-asparaginase II on T cells is a good
264 example of how bacterial subversion can lead to insight into basic host cell biology. In this
265 case these studies highlight the importance of asparagine as a nutrient in T cell metabolism
266 and activation (60).

267

268 Interference in lymphocyte chemotaxis

269 The majority of T cell-interacting bacterial proteins appear to mainly exert effects on
270 T cell activation, however, another strategy is to interfere with lymphocyte migration. For
271 example, *Shigella* exhibit the ability to directly invade T cells and cause an inhibition in their
272 chemokine-induced migration (61). *S. flexneri* are able to directly invade PMA-activated
273 CD4+ T cells, but not unstimulated, unactivated, primary CD4+ cells, with substantially
274 reduced responses to the chemoattractant CXCL12. CXCL12 signals through the chemokine
275 receptor CXCR4, the expression of which was not perturbed in these experiments (61). The
276 bacterial protein, IpgD, which can be secreted through the type III secretion apparatus (62),
277 has been implicated as being responsible for this activity, by acting on the pool of
278 intracellular phosphatidylinositol 4,5-bisphosphate (PIP2). Additionally, it would appear
279 that IpgD is able to act intracellularly in the absence of any other bacterial effectors (61).
280 These observations have been verified experimentally *in vivo* in mice, revealing that *S.*
281 *flexneri* target CD4+ T cells in the lymph node and confirming that invasion and migration
282 arrest occur *in vivo* (63). This discrimination between activated and non-activated T cells
283 could result in more specific targeting of activated T cells in the lamina propria rather than
284 the lymphoid follicles in the intestinal mucosa, thus targeting those cells that might actively
285 respond to infection. Further, a recent publication reported the ability of *Shigella* to inject
286 effectors into T cells in the absence of subsequent invasion, and suggest that the majority of
287 T cells are targeted by injection only, raising the possibility that the bacteria could use a “hit
288 and run” strategy to affect lymphocytes (64).

289

290 **Elimination of T cells**

291 A further strategy that is shared with more than one bacterial genus is seen with the
292 induction of apoptosis in T cells by the heat labile toxins, expressed by *E. coli* and *Vibrio*
293 *cholerae*, although they appear to have slightly different specificity and mode of action
294 between families and variants of the toxin. The heat labile toxins are structurally related
295 bacterial toxins that induce diarrhea in humans and animals (65). These toxins are
296 oligomers consisting of an A polypeptide bound to a pentameric array of B polypeptides.
297 The toxic effects are determined by the cell surface binding specificity of the B pentamers,
298 and the ADP ribosylating specificity of the A subunit (66). Cholera toxin (CT) produced by *V.*
299 *cholerae* binds to the ganglioside GM1 on epithelial cells via its B subunits, and when it is
300 trafficked to the cell cytosol, it catalyzes ADP ribosylation of adenylate cyclase, leading to
301 increased intracellular cAMP causing water secretion and diarrhea (67). However, it has
302 additional effects on other cells, including T cells. It was demonstrated some time ago that
303 CT was able to induce apoptosis in CD8+ T cells, although at that time the implications
304 during infection were unclear (68). More recently, it was confirmed that CT was able to
305 decrease the numbers of CD8 cells, and that this was not due to either a downregulation of
306 cell surface receptors, or selective proliferation of other cell types (69). Similarly, LTIIa from
307 *E. coli* is also able to deplete CD8+ T cells, likely by induction of apoptosis via cross-linking of
308 the ganglioside receptors, although this has not yet been explicitly demonstrated (69). In
309 mice injected with LT, transient induction of apoptosis mediated by glucocorticoids was
310 seen in all thymocyte subsets, although immature T cells were more affected than mature
311 cells (70). This effect was dependent on route of administration, and demonstrated that *in*
312 *vitro* treatment of cells did not entirely reflect the *in vivo* effects observed (70). In addition,
313 the maturation state of the T cell appears to determine the mechanism of apoptosis
314 triggered (71). Further, although CT does not appear to invoke apoptosis in CD4+ T cells, it

315 does appear to be able to inhibit activation, at least based on measurement of cell surface
316 expressed molecules (69).

317 There has been significant interest in using CT and LT as adjuvants in vaccination, and
318 understanding how it is able to steer T cell responses provides insight in how to better
319 improve vaccination, or to engineer non-toxic derivatives that are able to promote its
320 adjuvancy (72–75).

321 To our knowledge, there is only one other protein from a gastrointestinal bacterial
322 pathogen reported to have the ability to invoke apoptosis in T cells. This is the YpkA protein
323 of *Yersinia*, which is a multidomain protein with kinase activity. Expression of YpkA from a
324 mammalian expression vector transfected in Jurkat T cells induced significant apoptosis (76),
325 however, its role during infection is unclear.

326

327 **Concluding remarks**

328 It is evident from the examples above that gastrointestinal bacterial pathogens have
329 evolved diverse strategies to modulate lymphocyte function. However, the biological
330 significance of such activity during infection remains challenging to dissect, particularly for
331 factors that play additional roles in colonization. For such factors, the T lymphocyte
332 response to infection by a null mutant relative to the isogenic parent will be affected by the
333 magnitude and duration of exposure to bacterial antigens. One strategy to overcome this is
334 to use ligated intestinal loop models and recover intraepithelial lymphocytes exposed to
335 bacterial strains or their products *in situ* (e.g. (77)). Although, it can be challenging to
336 stimulate such cells to proliferate *ex vivo* and loop models often hold large numbers of
337 laboratory-cultured bacteria over the mucosa for a limited time, and thus do not simulate
338 the normal progression of gastrointestinal infection. It is noteworthy that attenuation of

339 mutants lacking some lymphocyte inhibitory factors is detected before one may anticipate
340 that adaptive responses have been generated, and further research is needed to understand
341 their impact on early pro-inflammatory responses and lymphocyte migration *in vivo*.
342 Further, for many of these proteins, little is known about their effects on T cells of specific
343 subsets and differentiation states. Knowledge of which might provide further insight into
344 their impact and timing of action during infection.

345 While some of the strategies outlined here rely on direct contact between the
346 pathogen and lymphocytes (e.g. via Type III secretion), in many cases inhibition relies on
347 diffusion of soluble proteins to meet their target cell type. Some of the factors described are
348 active in extremely low concentrations (e.g. lymphostatin acts in the femtomolar range;
349 (16)) and the extent to which lymphocytes in circulation are affected requires study. It is
350 evident from the ability of Shiga toxins to cause endothelial damage in kidney glomeruli that
351 proteins produced by gastrointestinal pathogens in the gut can act distally.

352 While the molecular basis of the activity of some lymphocyte inhibitory factors is
353 well understood (e.g. VacA, YopH, IpgD), for others a need exists to identify their cellular
354 targets and how their modification produces the observed phenotype. Such studies have
355 the potential to yield novel insights into both the basis of pathogenesis, but also the cellular
356 pathways and factors governing lymphocyte activation and function. With an understanding
357 of the mode of action of inhibitory factors, it may also become feasible to design new
358 treatments. For example, with the knowledge that *Helicobacter* may use γ -
359 glutamyltranspeptidase to restrict lymphocyte activation via interference in glutamate
360 metabolism, researchers have recently demonstrated that oral glutathione supplementation

361 can reduce gastric pathology and inflammation due to *H. suis* in gerbils (78). The extent to
362 which this is a consequence of altered T lymphocyte function requires further study.

363 It is striking that relatively little direct duplication of strategies to inhibit lymphocyte
364 function has been identified across bacterial genera. Nevertheless, the vast quantities of
365 sequence data now generated for pathogens will facilitate the identification of homologs of
366 lymphocyte inhibitory factors that may be relevant in other diseases and differ in
367 mechanism. For example, a family of proteins homologous to lymphostatin occur in diverse
368 *Chlamydia* species of veterinary and public health importance and share predicted
369 glycosyltransferase motifs (79).

370 In addition to evaluating the value of lymphocyte inhibitory factors as subunit
371 vaccines or as targets for novel inhibitors, merit exists in exploring the therapeutic potential
372 of such molecules for disorders associated with lymphocyte proliferation or activity. A
373 challenge of such will be ensuring specific targeting of pathology-associated lymphocytes
374 without deleterious effects on immune function.

375

376 ACKNOWLEDGEMENTS

377 The authors gratefully acknowledge strategic investment from the Biotechnology and
378 Biological Sciences Research Council (references BB/J004227/1 and BB/P013740/1).

379

380 FIGURES

381 FIGURE 1: GALT and T cell distribution in the intestine

382 Immune responses in the intestine are mainly controlled by gut associated lymphoid tissue
383 (GALT), including the Peyer patches, mesenteric lymph nodes and isolated lymphoid
384 follicles in the mucosa and lamina propria. The mucosa is also studded with intraepithelial
385 lymphocytes. Naïve T cells can be recruited from the circulation to lymphoid organs in the
386 intestines, where they can be activated.

387 FIGURE 2: Summary of T cell interacting bacterial proteins and their targets

388 The majority of bacterial proteins that interact with T cells are directed at modifying
389 activation/proliferation, however, there are some proteins that affect chemotaxis and
390 apoptosis. Where the key affected molecules are known, these are indicated, however, the
391 details of a number of molecules remain unknown. Bacterial protein names are bounded by
392 gray boxes. sAg= superantigen (*Staphylococcus*), CT= cholera toxin(*Vibrio cholera*),
393 GGT=gamma glutamyl transferase (*H. pylori*, *C. jejuni*), lpgD= invasion plasmid gene D
394 (*Shigella*), LifA= lymphocyte inhibiting factor A (*E. coli*), LT= heat labile toxin (*E. coli*),
395 STM3106=asparaginase (*Salmonella*), APC= antigen presenting cell, MHC= major
396 histocompatibility complex.

397 REFERENCES

- 398 1. Kirk MD, Pires SM, Black RE, Caipo M, Crump JA, Devleesschauwer B, Döpfer D, Fazil
399 A, Fischer-Walker CL, Hald T, Hall AJ, Keddy KH, Lake RJ, Lanata CF, Torgerson PR,
400 Havelaar AH, Angulo FJ. 2015. World Health Organization estimates of the global and
401 regional disease burden of 22 foodborne bacterial, protozoal, and viral diseases,
402 2010: A Data synthesis. PLoS Med **12**:e1001921.
- 403 2. Santos AS, Finlay BB. 2015. Bringing down the host: Enteropathogenic and

- 404 enterohaemorrhagic *Escherichia coli* effector-mediated subversion of host innate
405 immune pathways. *Cell Microbiol* **17**:318–332.
- 406 3. **Deng W, Marshall NC, Rowland JL, McCoy JM, Worrall LJ, Santos AS, Strynadka NCJ,**
407 **Finlay BB.** 2017. Assembly, structure, function and regulation of type III secretion
408 systems. *Nat Rev Microbiol* **15**:323–337.
- 409 4. **van Wijk F, Cheroutre H.** 2010. Mucosal T cells in gut homeostasis and inflammation.
410 *Expert Rev Clin Immunol* **6**:559–66.
- 411 5. **Koboziev I, Karlsson F, Grisham MB.** 2010. Gut-associated lymphoid tissue, T cell
412 trafficking, and chronic intestinal inflammation. *Ann N Y Acad Sci* **E86**–93.
- 413 6. **Barber MA.** 1914. Milk poisoning due to a type of *Staphylococcus albus* occurring in
414 the udder of a healthy cow. *Philipp J Sci* **9B**:515.
- 415 7. **Choi YW, Kotzin B, Herron L, Callahan J, Marrack P, Kappler J.** 1989. Interaction of
416 *Staphylococcus aureus* toxin superantigens with human T cells. *Proc Natl Acad Sci U S*
417 *A* **86**:8941–5.
- 418 8. **Krakauer T, Pradhan K, Stiles BG.** 2016. Staphylococcal superantigens spark host-
419 mediated danger signals. *Front Immunol* **7**:Article 23.
- 420 9. **Krakauer T.** 2013. Update on staphylococcal superantigen-induced signaling
421 pathways and therapeutic interventions. *Toxins (Basel)* **5**:1629–54.
- 422 10. **Stuart PM, Woodward JG.** 1992. *Yersinia enterocolitica* produces superantigenic
423 activity. *J Immunol* **148**:225–33.
- 424 11. **Abe J, Takeda T, Watanabe Y, Nakao H, Kobayashi N, Leung DY, Kohsaka T.** 1993.

- 425 Evidence for superantigen production by *Yersinia pseudotuberculosis*. J Immunol
426 **151**:4183–8.
- 427 12. **Amphlett A.** 2016. Far East Scarlet-Like Fever: A review of the epidemiology,
428 symptomatology, and role of superantigenic toxin: *Yersinia pseudotuberculosis*-
429 derived mitogen A. Open forum Infect Dis **3**:ofv202.
- 430 13. **Carnoy C, Mullet C, Müller-Alouf H, Leteurtre E, Simonet M.** 2000. Superantigen
431 YPMa exacerbates the virulence of *Yersinia pseudotuberculosis* in mice. Infect Immun
432 **68**:2553–9.
- 433 14. **Goubard A, Loïez C, Abe J, Fichel C, Herwegh S, Faveeuw C, Porte R, Cayet D,**
434 **Sebbane F, Penet S, Foligné B, Desreumaux P, Saito H, Sirard J-C, Simonet M, Carnoy**
435 **C.** 2015. Superantigenic *Yersinia pseudotuberculosis* induces the expression of
436 granzymes and perforin by CD4 + T Cells. Infect Immun **83**:2053–2064.
- 437 15. **Klapproth JM, Scaletsky IC, McNamara BP, Lai LC, Malstrom C, James SP,**
438 **Donnenberg MS.** 2000. A large toxin from pathogenic *Escherichia coli* strains that
439 inhibits lymphocyte activation. Infect Immun **68**:2148–55.
- 440 16. **Cassady-Cain RL, Blackburn EA, Alsarraf H, Dedic E, Bease AG, Böttcher B, Jørgensen**
441 **R, Wear M, Stevens MP.** 2016. Biophysical characterization and activity of
442 lymphostatin, a multifunctional virulence factor of attaching and effacing *Escherichia*
443 *coli*. J Biol Chem **291**:5803–16.
- 444 17. **Klapproth JM, Donnenberg MS, Abraham JM, Mobley HL, James SP.** 1995. Products
445 of enteropathogenic *Escherichia coli* inhibit lymphocyte activation and lymphokine
446 production. Infect Immun **63**:2248–54.

- 447 18. **Cassady-Cain RL, Blackburn EA, Bell CR, Elshina E, Hope JC, Stevens MP.** 2017.
448 Inhibition of antigen-specific and nonspecific stimulation of bovine T and B cells by
449 lymphostatin from attaching and effacing *Escherichia coli*. *Infect Immun* **85**:e00845–
450 16.
- 451 19. **Stevens MP, van Diemen PM, Frankel G, Phillips AD, Wallis TS.** 2002. Efa1 influences
452 colonization of the bovine intestine by shiga toxin-producing *Escherichia coli*
453 serotypes O5 and O111. *Infect Immun* **70**:5158–66.
- 454 20. **Deacon V, Dziva F, van Diemen PM, Frankel G, Stevens MP.** 2010. Efa-1/LifA
455 mediates intestinal colonization of calves by enterohaemorrhagic *Escherichia coli*
456 O26 : H- in a manner independent of glycosyltransferase and cysteine protease motifs
457 or effects on type III secretion. *Microbiology* **156**:2527–36.
- 458 21. **Klapproth J-MA, Sasaki M, Sherman M, Babbin B, Donnenberg MS, Fernandes PJ,**
459 **Scaletsky ICA, Kalman D, Nusrat A, Williams IR.** 2005. *Citrobacter rodentium* *lifA/efa1*
460 is essential for colonic colonization and crypt cell hyperplasia in vivo. *Infect Immun*
461 **73**:1441–51.
- 462 22. **Nicholls L, Grant TH, Robins-Browne RM.** 2000. Identification of a novel genetic locus
463 that is required for *in vitro* adhesion of a clinical isolate of enterohaemorrhagic
464 *Escherichia coli* to epithelial cells. *Mol Microbiol* **35**:275–88.
- 465 23. **Boncristiano M, Paccani SR, Barone S, Ulivieri C, Patrussi L, Ilver D, Amedei A,**
466 **D’Elios MM, Telford JL, Baldari CT.** 2003. The *Helicobacter pylori* vacuolating toxin
467 inhibits T cell activation by two independent mechanisms. *J Exp Med* **198**:1887–97.
- 468 24. **Gebert B, Fischer W, Weiss E, Hoffmann R, Haas R.** 2003. *Helicobacter pylori*

- 469 vacuolating cytotoxin inhibits T lymphocyte activation. *Science* (80-) **301**:1099–1102.
- 470 25. **Sundrud MS, Torres VJ, Unutmaz D, Cover TL.** 2004. Inhibition of primary human T
471 cell proliferation by *Helicobacter pylori* vacuolating toxin (VacA) is independent of
472 VacA effects on IL-2 secretion. *Proc Natl Acad Sci* **101**:7727–7732.
- 473 26. **Utsch C, Haas R.** 2016. VacA's induction of VacA-containing vacuoles (VCVs) and their
474 immunomodulatory activities on human T cells. *Toxins (Basel)* **8**:190.
- 475 27. **Foegeding NJ, Caston RR, McClain MS, Ohi MD, Cover TL.** 2016. An overview of
476 *Helicobacter pylori* VacA toxin biology. *Toxins (Basel)* **8**:173.
- 477 28. **Atherton JC, Cao P, Peek RM, Tummuru MK, Blaser MJ, Cover TL.** 1995. Mosaicism in
478 vacuolating cytotoxin alleles of *Helicobacter pylori*. Association of specific vacA types
479 with cytotoxin production and peptic ulceration. *J Biol Chem* **270**:17771–7.
- 480 29. **Torres VJ, Ivie SE, McClain MS, Cover TL.** 2005. Functional properties of the p33 and
481 p55 domains of the *Helicobacter pylori* vacuolating cytotoxin. *J Biol Chem* **280**:21107–
482 21114.
- 483 30. **Sewald X, Gebert-Vogl B, Prassl S, Barwig I, Weiss E, Fabbri M, Osicka R, Schiemann**
484 **M, Busch DH, Semmrich M, Holzmann B, Sebo P, Haas R.** 2008. Integrin subunit CD18
485 is the T-lymphocyte receptor for the *Helicobacter pylori* vacuolating cytotoxin. *Cell*
486 *Host Microbe* **3**:20–29.
- 487 31. **Sewald X, Jiménez-Soto L, Haas R.** 2011. PKC-dependent endocytosis of the
488 *Helicobacter pylori* vacuolating cytotoxin in primary T lymphocytes. *Cell Microbiol*
489 **13**:482–496.
- 490 32. **Kern B, Jain U, Utsch C, Otto A, Busch B, Jiménez-Soto L, Becher D, Haas R.** 2015.

- 491 Characterization of *Helicobacter pylori* VacA-containing vacuoles (VCVs), VacA
492 intracellular trafficking and interference with calcium signalling in T lymphocytes. *Cell*
493 *Microbiol* **17**:1811–1832.
- 494 33. **Salama NR, Otto G, Tompkins L, Falkow S.** 2001. Vacuolating cytotoxin of
495 *Helicobacter pylori* plays a role during colonization in a mouse model of infection.
496 *Infect Immun* **69**:730–6.
- 497 34. **Cornelis GR, Biot T, Rouvroit CL, Michiels T, Mulder B, Sluiter C, Sory M-P,**
498 **Bouchaute M, Vanooteghem J-C.** 1989. The *Yersinia* yop regulon. *Mol Microbiol*
499 **3**:1455–1459.
- 500 35. **Guan KL, Dixon JE.** 1990. Protein tyrosine phosphatase activity of an essential
501 virulence determinant in *Yersinia*. *Science* **249**:553–6.
- 502 36. **Yao T, Mecsas J, Healy JJ, Falkow S, Chien Y.** 1999. Suppression of T and B
503 lymphocyte activation by a *Yersinia pseudotuberculosis* virulence factor, yopH. *J Exp*
504 *Med* **190**:1343–50.
- 505 37. **Sauvonnet N, Lambermont I, Bruggen P van der, Cornelis GR.** 2002. YopH prevents
506 monocyte chemoattractant protein 1 expression in macrophages and T-cell
507 proliferation through inactivation of the phosphatidylinositol 3-kinase pathway. *Mol*
508 *Microbiol* **45**:805–815.
- 509 38. **Alonso A, Bottini N, Bruckner S, Rahmouni S, Williams S, Schoenberger SP, Mustelin**
510 **T.** 2004. Lck dephosphorylation at Tyr-394 and inhibition of T Cell antigen receptor
511 signaling by *Yersinia* phosphatase YopH. *J Biol Chem* **279**:4922–4928.
- 512 39. **de la Puerta ML, Trinidad AG, Rodríguez M del C, Bogetz J, Sánchez Crespo M,**

- 513 **Mustelin T, Alonso A, Bayón Y.** 2009. Characterization of new substrates targeted by
514 *Yersinia* tyrosine phosphatase YopH. PLoS One **4**:e4431.
- 515 40. **Trülsch K, Sporleder T, Igwe EI, Rüssmann H, Heesemann J.** 2004. Contribution of
516 the major secreted yops of *Yersinia enterocolitica* O:8 to pathogenicity in the mouse
517 infection model. Infect Immun **72**:5227–34.
- 518 41. **Fisher ML, Castillo C, Mecsas J.** 2007. Intranasal inoculation of mice with *Yersinia*
519 *pseudotuberculosis* causes a lethal lung infection that is dependent on *Yersinia* outer
520 proteins and PhoP. Infect Immun **75**:429–42.
- 521 42. **Arencibia I, Suárez NC, Wolf-Watz H, Sundqvist KG.** 1997. *Yersinia* invasin, a bacterial
522 beta1-integrin ligand, is a potent inducer of lymphocyte motility and migration to
523 collagen type IV and fibronectin. J Immunol **159**:1853–9.
- 524 43. **Leo JC, Oberhettinger P, Schütz M, Linke D.** 2015. The inverse autotransporter family:
525 Intimin, invasin and related proteins. Int J Med Microbiol **305**:276–282.
- 526 44. **Gonçalves NS, Hale C, Dougan G, Frankel G, MacDonald TT.** 2003. Binding of intimin
527 from enteropathogenic *Escherichia coli* to lymphocytes and its functional
528 consequences. Infect Immun **71**:2960–5.
- 529 45. **Higgins LM, Frankel G, Connerton I, Gonçalves NS, Dougan G, MacDonald TT.** 1999.
530 Role of bacterial intimin in colonic hyperplasia and inflammation. Science **285**:588–
531 91.
- 532 46. **Patel CH, Powell JD.** 2017. Targeting T cell metabolism to regulate T cell activation,
533 differentiation and function in disease. Curr Opin Immunol **46**:82–88.
- 534 47. **MacIver NJ, Michalek RD, Rathmell JC.** 2013. Metabolic regulation of T lymphocytes.

- 535 Annu Rev Immunol **31**:259–83.
- 536 48. **Floch P, Pey V, Castroviejo M, Dupuy JW, Bonneu M, de la Guardia AH, Pitard V,**
537 **Mégraud F, Lehours P.** 2014. Role of *Campylobacter jejuni* gamma-glutamyl
538 transpeptidase on epithelial cell apoptosis and lymphocyte proliferation. Gut Pathog
539 **6**:20.
- 540 49. **Barnes IHA, Bagnall MC, Browning DD, Thompson SA, Manning G, Newell DG.** 2007.
541 Gamma-glutamyl transpeptidase has a role in the persistent colonization of the avian
542 gut by *Campylobacter jejuni*. Microb Pathog **43**:198–207.
- 543 50. **Chevalier C, Thiberge JM, Ferrero RL, Labigne A.** 1999. Essential role of *Helicobacter*
544 *pylori* gamma-glutamyltranspeptidase for the colonization of the gastric mucosa of
545 mice. Mol Microbiol **31**:1359–72.
- 546 51. **Rossi M, Bolz C, Revez J, Javed S, El-Najjar N, Anderl F, Hyytiäinen H, Vuorela P,**
547 **Gerhard M, Hänninen M-L.** 2012. Evidence for conserved function of γ -
548 glutamyltranspeptidase in *Helicobacter* Genus. PLoS One **7**:e30543.
- 549 52. **Zhang G, Ducatelle R, Pasmans F, D’Herde K, Na, Huang L, Smet A, Haesebrouck F,**
550 **Flahou B.** 2013. Effects of *Helicobacter suis* γ -glutamyl transpeptidase on
551 lymphocytes: modulation by glutamine and glutathione supplementation and outer
552 membrane vesicles as a putative delivery route of the enzyme. PLoS One **8**:e77966.
- 553 53. **Schmees C, Prinz C, Treptau T, Rad R, Hengst L, Volland P, Bauer S, Brenner L, Schmid**
554 **RM, Gerhard M.** 2007. Inhibition of T-cell proliferation by *Helicobacter pylori* γ -
555 glutamyl transpeptidase. Gastroenterology **132**:1820–1833.
- 556 54. **Wüstner S, Mejías-Luque R, Koch MF, Rath E, Vieth M, Sieber SA, Haller D, Gerhard**

- 557 M. 2015. *Helicobacter pylori* γ -glutamyltranspeptidase impairs T-lymphocyte function
558 by compromising metabolic adaption through inhibition of cMyc and IRF4 expression.
559 Cell Microbiol **17**:51–61.
- 560 55. van der Velden AWM, Copass MK, Starnbach MN. 2005. *Salmonella* inhibit T cell
561 proliferation by a direct, contact-dependent immunosuppressive effect. Proc Natl
562 Acad Sci U S A **102**:17769–74.
- 563 56. McLaughlin PA, McClelland M, Yang H-J, Porwollik S, Bogomolnaya L, Chen J-S,
564 Andrews-Polymenis H, van der Velden AWM. 2017. Contribution of Asparagine
565 Catabolism to *Salmonella* Virulence. Infect Immun **85**:e00740–16.
- 566 57. van der Velden AWM, Dougherty JT, Starnbach MN. 2008. Down-modulation of TCR
567 expression by *Salmonella enterica* serovar Typhimurium. J Immunol **180**:5569–74.
- 568 58. Kullas AL, McClelland M, Yang H-J, Tam JW, Torres A, Porwollik S, Mena P, McPhee
569 JB, Bogomolnaya L, Andrews-Polymenis H, van der Velden AWM. 2012. L-
570 asparaginase II produced by *Salmonella typhimurium* inhibits T cell responses and
571 mediates virulence. Cell Host Microbe **12**:791–8.
- 572 59. Chaudhuri RR, Morgan E, Peters SE, Pleasance SJ, Hudson DL, Davies HM, Wang J,
573 van Diemen PM, Buckley AM, Bowen AJ, Pullinger GD, Turner DJ, Langridge GC,
574 Turner AK, Parkhill J, Charles IG, Maskell DJ, Stevens MP. 2013. Comprehensive
575 assignment of roles for *Salmonella* Typhimurium genes in intestinal colonization of
576 food-producing animals. PLoS Genet **9**:e1003456.
- 577 60. Torres A, Luke JD, Kullas AL, Kapilashrami K, Botbol Y, Koller A, Tonge PJ, Chen EI,
578 Macian F, van der Velden AWM. 2016. Asparagine deprivation mediated by

- 579 *Salmonella* asparaginase causes suppression of activation-induced T cell metabolic
580 reprogramming. *J Leukoc Biol* **99**:387–98.
- 581 61. **Konradt C, Frigimelica E, Nothelfer K, Puhar A, Salgado-Pabon W, di Bartolo V, Scott-**
582 **Algara D, Rodrigues CD, Sansonetti PJ, Phalipon A.** 2011. The *Shigella flexneri* Type
583 three secretion system effector IpgD inhibits T cell migration by manipulating host
584 phosphoinositide metabolism. *Cell Host Microbe* **9**:263–272.
- 585 62. **Parsot C.** 2009. *Shigella* type III secretion effectors: how, where, when, for what
586 purposes? *Curr Opin Microbiol* **12**:110–116.
- 587 63. **Salgado-Pabón W, Celli S, Arena ET, Nothelfer K, Roux P, Sellge G, Frigimelica E,**
588 **Bousso P, Sansonetti PJ, Phalipon A.** 2013. *Shigella* impairs T lymphocyte dynamics *in*
589 *vivo*. *Proc Natl Acad Sci U S A* **110**:4458–63.
- 590 64. **Pinaud L, Samassa F, Porat Z, Ferrari ML, Belotserkovsky I, Parsot C, Sansonetti PJ,**
591 **Campbell-Valois F-X, Phalipon A.** 2017. Injection of T3SS effectors not resulting in
592 invasion is the main targeting mechanism of *Shigella* toward human lymphocytes.
593 *Proc Natl Acad Sci U S A* **114**:9954–9959.
- 594 65. **Gill DM, Clements JD, Robertson DC, Finkelstein RA.** 1981. Subunit number and
595 arrangement in *Escherichia coli* heat-labile enterotoxin. *Infect Immun* **33**:677–82.
- 596 66. **Spangler BD.** 1992. Structure and function of cholera toxin and the related
597 *Escherichia coli* heat-labile enterotoxin. *Microbiol Rev* **56**:622–47.
- 598 67. **Baldauf K, Royal J, Hamorsky K, Matoba N.** 2015. Cholera toxin B: One subunit with
599 many pharmaceutical applications. *Toxins (Basel)* **7**:974–996.
- 600 68. **Yankelevich B, Soldatenkov VA, Hodgson J, Polotsky AJ, Creswell K, Mazumder A.**

1996. Differential induction of programmed cell death in CD8+and CD4+T cells by the B subunit of cholera toxin. *Cell Immunol* **168**:229–234.
69. **Arce S, Nawar HF, Russell MW, Connell TD.** 2005. Differential binding of *Escherichia coli* enterotoxins LT-IIa and LT-IIb and of cholera toxin elicits differences in apoptosis, proliferation, and activation of lymphoid cells. *Infect Immun* **73**:2718–27.
70. **Tamayo E, Merino R, González-Rojas J, Marquina R, Santiuste I, Amado JA, Rappuoli R, Del Giudice G, Merino J.** 2005. The *Escherichia coli* heat-labile enterotoxin induces apoptosis of immature lymphocytes in vivo via a glucocorticoid-dependent pathway. *Eur J Immunol* **35**:3505–3515.
71. **Tamayo E, Postigo J, Del Giudice G, Rappuoli R, Benito A, Yagita H, Merino R, Merino J.** 2009. Involvement of the intrinsic and extrinsic cell-death pathways in the induction of apoptosis of mature lymphocytes by the *Escherichia coli* heat-labile enterotoxin. *Eur J Immunol* **39**:439–446.
72. **Basset C, Thiam F, Di Martino C, Holton J, Clements JD, Kohli E.** 2010. Cholera-like enterotoxins and regulatory T cells. *Toxins (Basel)* **2**:1774–1795.
73. **Tsai H-C, Wu R.** 2015. Mechanisms of cholera toxin in the modulation of TH17 responses. *Crit Rev Immunol* **35**:135–52.
74. **Storset AK, Kulberg S, Berg I, Boysen P, Hope JC, Dissen E.** 2004. NKp46 defines a subset of bovine leukocytes with natural killer cell characteristics. *Eur J Immunol* **34**:669–676.
75. **Lebens M, Terrinoni M, Karlsson SL, Larena M, Gustafsson-Hedberg T, Källgård S, Nygren E, Holmgren J.** 2016. Construction and preclinical evaluation of mmCT, a

- 623 novel mutant cholera toxin adjuvant that can be efficiently produced in genetically
624 manipulated *Vibrio cholerae*. *Vaccine* **34**:2121–2128.
- 625 76. **Park H, Teja K, O'Shea JJ, Siegel RM.** 2007. The *Yersinia* effector protein YpkA induces
626 apoptosis independently of actin depolymerization. *J Immunol* **178**:6426–34.
- 627 77. **Menge C, Blessenohl M, Eisenberg T, Stamm I, Baljer G.** 2004. Bovine ileal
628 intraepithelial lymphocytes represent target cells for Shiga toxin 1 from *Escherichia*
629 *coli*. *Infect Immun* **72**:1896–905.
- 630 78. **De Bruyne E, Ducatelle R, Foss D, Sanchez M, Joosten M, Zhang G, Smet A, Pasmans**
631 **F, Haesebrouck F, Flahou B.** 2016. Oral glutathione supplementation drastically
632 reduces *Helicobacter*-induced gastric pathologies. *Sci Rep* **6**:20169.
- 633 79. **Belland RJ, Scidmore MA, Crane DD, Hogan DM, Whitmire W, McClarty G, Caldwell**
634 **HD.** 2001. *Chlamydia trachomatis* cytotoxicity associated with complete and partial
635 cytotoxin genes. *Proc Natl Acad Sci U S A* **98**:13984–9.
- 636



